

Integrability of anisotropic and homogeneous Universes in scalar-tensor theory of gravitation

Julien Larena

Laboratoire Univers et Théories, UMR8102, Paris Observatory and University Paris 7 Denis Diderot, julien.larena@obspm.fr

Jérôme Perez

Laboratoire de Mathématiques Appliquées, Ecole Nationale Supérieure de Techniques Avancées, jerome.perez@ensta.fr

Abstract. In this paper, we develop a method based on the analysis of the Kovalewski exponents to study the integrability of anisotropic and homogeneous Universes. The formalism is developed in scalar-tensor gravity, the general relativistic case appearing as a special case of this larger framework. Then, depending on the rationality of the Kovalewski exponents, the different models, both in the vacuum and in presence of a barotropic matter fluid, are classified, and their integrability is discussed.

Greek indexes run from 0 to 4, Latin indexes run from 1 to 3 and ∇^μ indicates a covariant derivative.

PACS numbers: 02.30.Ik, 04.20.-q, 04.20.Cv, 98.80.Jk

1. Dynamical equations

1.1. Homogeneous spaces

The classification of homogeneous and anisotropic $3D$ spaces gives 11 different types of spaces associated to distinct families of structure group constants

$$C_{ab}^c = \varepsilon_{abd} N^{dc} + \delta_b^c A_a - \delta_a^c A_b \quad (1)$$

where ε_{abd} is the usual totally antisymmetric unit tensor, δ_b^c the Kronecker symbol, N^{ab} is the contravariant component of an order 2 symmetric tensor, and the vector A must follows

$$N^{ab}A_b = N_{ab}A^b = 0. \quad (2)$$

Without loss generality, one can write $N^{ab} = \text{diag}(n_1, n_2, n_3)$ and $A_b = [a, 0, 0]$ provided that $an_1 = 0$. Distinct homogeneous and anisotropic spaces in $3D$ can then be classified in the following table:

	n_1	n_2	n_3	a	Name
0 is eigenvalue of N with multiplicity 3	0	0	0	0	B_I
	0	0	0	$\forall$	B_V
0 is eigenvalue of N with multiplicity 2	1	0	0	0	B_{II}
	0	1	0	$\forall$	B_{IV}
0 is eigenvalue of N with multiplicity 1	1	1	0	0	B_{VII_o}
	0	1	1	$\forall$	B_{VII_a}
	1	-1	0	0	B_{VI_o}
	0	1	-1	$\neq 1$	B_{VI_a}
	0	1	-1	1	B_{III}
0 is not an eigenvalue of N	1	1	1	0	B_{IX}
	1	1	-1	0	B_{VIII}

Note that each case is degenerate, for example $n_1 = -1, n_2 = 1, n_3 = -1$ and $a = 0$ is in the equivalence class of B_{IX} which contains all the possibilities with a positive signature and 0 out of the spectrum of N . This is the well known Bianchi classification (see [1],[2]).

1.2. Scalar-tensor theory of gravitation

In scalar-tensor theories of gravity, the dynamics of the Universe contains a new scalar degree of freedom that couples explicitly to the energy content of the Universe [3, 4, 5, 6, 7]. In units of $c = 1$, the action generically writes, in the so-called Einstein frame:

$$S = \frac{1}{4\pi G} \int \left(\frac{R}{4} - \frac{1}{2} \varphi_{,\mu} \varphi^{,\mu} - U(\varphi) \right) \sqrt{-g} d^4x \\ + S_m(\psi_m, \Theta^2(\varphi) g_{\mu\nu}), \quad (3)$$

G being a bare gravitational constant, φ the scalar field, $U(\varphi)$ its self-interaction term and $\Theta(\varphi)$ its coupling to matter. The functional $S_m(\psi_m, \Theta^2(\varphi)g_{\mu\nu})$ stands for the action of any field ψ_m that contributes to the energy content of the Universe. It expresses the fact that all these fields couple universally to a conformal metric $\tilde{g}_{\mu\nu} = \Theta^2(\varphi)g_{\mu\nu}$, then implying that the weak equivalence principle (local universality of free fall for non-gravitationally bound objects) holds in this class of theories. The metric $\tilde{g}_{\mu\nu}$ defines the Dicke-Jordan frame, in which standard rods and clocks can be used to make measurements (since in this frame, the matter part of the action acquires its standard form). Despite the conformal relation, these two frames have a different status: in the Dicke-Jordan frame, where the gravitational degrees of freedom are mixed, the Lagrangian for the matter fields does not contain explicitly the new scalar field: the non gravitational physics has then its standard form. In the Einstein frame, the scalar degree of freedom explicitly couples to the matter fields, then leading for example to the variation of the inertial masses of point-like particles. Of course, the two frames describe the same physical world. Nevertheless, the usual interpretation of the observable quantities is profoundly modified in the Einstein frame, whereas it holds in the Dicke-Jordan frame, where the rods and clocks made with matter are not affected by the presence of the scalar field. That is why one usually refers to the Dicke-Jordan frame as the observable one. However, the dynamics of the fields is generally more easily described in the Einstein frame, so that in this work, since we are interested in the integrability of the models rather than in their physical content, the analysis will be done in the Einstein frame.

Varying the Einstein frame action (3) with respect to the fields yields the equations:

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \chi T_{\mu\nu} + T_{\mu\nu}^{\varphi} \quad (4)$$

$$\square\varphi = -\frac{\chi}{2}\omega(\varphi)T + \frac{dU(\varphi)}{d\varphi} \quad (5)$$

$$\nabla_{\nu}T_{\mu}^{\nu} = \omega(\varphi)T\nabla_{\mu}\varphi \quad (6)$$

where $\chi = 8\pi G$, T is the trace of the energy-momentum tensor of matter fields $T_{\mu\nu}$, and $T_{\mu\nu}^{\varphi} = 2\varphi_{,\mu}\varphi_{,\nu} - g_{\mu\nu}(g^{\alpha\beta}\varphi_{,\alpha}\varphi_{,\beta}) - 2U(\varphi)g_{\mu\nu}$ is the energy-momentum tensor of the scalar field. Moreover, we have defined the coupling $\omega(\varphi) = \frac{d\ln\Theta}{d\varphi}$. It is important to note that these equations reduce to those of General Relativity in presence of a scalar field iff $\omega(\varphi) = 0$.

1.3. Dynamical equations of scalar-tensor theory in homogeneous spaces

If one denotes by t the physical time and by τ a conformal time such that $dt = N(\tau)d\tau$, where $N(\tau)$ is generally called "lapse function", the line element of the 3 + 1 physical space reads:

$$ds^2 = g_{ij}dx^i dx^j - dt^2 = \gamma(\tau)\omega^i\omega^j - N^2(\tau)d\tau^2 \quad (7)$$

As explained in [8], one can find an invariant basis of differential forms $\{\omega^1, \omega^2, \omega^3\}$ in each Bianchi space such that γ is diagonal; we note hereafter $\gamma(.) = \text{diag}(\alpha_i(.))$.

Taking $V(\tau) = (\alpha_1 \alpha_2 \alpha_3)^{1/2}$ as a lapse function, and introducing $A_i = \ln(\alpha_i)$, the scalar-tensor dynamical equations in homogeneous and anisotropic Universe read:

$$\left\{ \begin{array}{lcl} \chi V^2 [3(P + P_\varphi) + \rho + \rho_\varphi] & = & -(\ln(V^2))'' + \frac{1}{2}((\ln(V^2))')^2 \\ & & -\frac{1}{2}(A_1'^2 + A_2'^2 + A_3'^2) \\ \chi V^2 [\rho + \rho_\varphi - (P + P_\varphi)] & = & A_1'' + n_1^2 \alpha_1^2 - (n_2 \alpha_2 - n_3 \alpha_3)^2 \\ \chi V^2 [\rho + \rho_\varphi - (P + P_\varphi)] & = & A_2'' + n_2^2 \alpha_2^2 - (n_3 \alpha_3 - n_1 \alpha_1)^2 \\ \chi V^2 [\rho + \rho_\varphi - (P + P_\varphi)] & = & A_3'' + n_3^2 \alpha_3^2 - (n_1 \alpha_1 - n_2 \alpha_2)^2 \\ \chi V^2 \omega(\varphi) [\rho - 3P] & = & -2\varphi'' - 4(\ln V)' \varphi' - 2 \frac{dU}{d\varphi} V^2 \end{array} \right. \quad (8)$$

In these equations we have written ' for $d/d\tau$; in addition the Universe is filled by a perfect fluid with pressure P and energy density ρ ; finally we have noted:

$$\left\{ \begin{array}{lcl} \rho_\varphi & := & [\varphi'^2/2 + U(\varphi)]/\chi \\ P_\varphi & := & [\varphi'^2/2 - U(\varphi)]/\chi \end{array} \right. \quad (9)$$

Taking $U \equiv 0$, and reorganizing the first dynamical equation using the three others, the system (8) becomes:

$$\left\{ \begin{array}{lcl} 0 & = & E_c + E_p - 4\chi\rho V^2 - 4\chi\rho_\varphi V^2 \\ \chi V^2 [\rho + \rho_\varphi - (P + P_\varphi)] & = & A_1'' + n_1^2 \alpha_1^2 - (n_2 \alpha_2 - n_3 \alpha_3)^2 \\ \chi V^2 [\rho + \rho_\varphi - (P + P_\varphi)] & = & A_2'' + n_2^2 \alpha_2^2 - (n_3 \alpha_3 - n_1 \alpha_1)^2 \\ \chi V^2 [\rho + \rho_\varphi - (P + P_\varphi)] & = & A_3'' + n_3^2 \alpha_3^2 - (n_1 \alpha_1 - n_2 \alpha_2)^2 \\ \chi V^2 \omega(\varphi) [\rho - 3P] & = & -2\varphi'' - 4(\ln V)' \varphi' \end{array} \right. \quad (10)$$

where

$$E_c := A_1' A_2' + A_1' A_3' + A_3' A_2' \quad \text{and} \quad E_p := \sum_{i \neq j=1}^3 n_i n_j e^{A_i + A_j} - \sum_{i=1}^3 n_i^2 e^{2A_i}$$

The energy-momentum conservation, that in scalar-tensor theory is:

$$\nabla^\mu T_{\mu\nu} = \omega(\varphi) T_{\lambda\beta} g^{\lambda\beta} \partial_\nu \varphi \quad (11)$$

coupled to the hypothesis of a barotropic fluid:

$$P = (\Gamma - 1) \rho \quad (12)$$

allows us to obtain a relation between ρ , V and φ which is:

$$\rho = \rho_o V^{-\Gamma} \Theta^{4-3\Gamma} \quad \text{with } \rho_o \in \mathbb{R}^+ \setminus \{0\} \quad (13)$$

One should note that a fluid of radiation ($\Gamma = 4/3$) doesn't couple directly to the scalar field. Assuming a power law dependence of φ' in V , more precisely:

$$\varphi' = \varphi'_o V^\Delta \quad (14)$$

and making use of (13) one can solve the last equation of system (8), and obtain an explicit dependence on V for $\Theta(\varphi)$:

$$\rho_o \Theta^{4-3\Gamma} = \frac{a^2}{2\chi} V^{2(\Delta-1)+\Gamma} \quad \text{where } a^2 = \frac{4(\Delta+2)}{2(1-\Delta)-\Gamma} \varphi_o'^2 \quad (15)$$

Assuming $\rho_o > 0$ (which corresponds to non exotic matter) and $0 \leq \Gamma \leq 2$ (which corresponds to the largest class of barotropic fluid), relation (15) holds under the condition:

$$-2 < \Delta < \frac{2 - \Gamma}{2} \quad (16)$$

The assumption (14) is a constraint on the entire dynamical system rather than on the scalar-tensor theory itself. Indeed, any choice of coupling function $\omega(\varphi)$ can be done, but then the resulting behavior of $V(\tau)$ is completely fixed by the last equation of system (10). Conversely, imposing a behavior for the volume $V(\tau)$ determines the corresponding coupling function. For example, imposing a Brans-Dicke theory, i.e. $\omega(\varphi) = \omega_0 = cste$ results in $V(\tau) \propto \tau^{1/(2-\Delta)}$; on the contrary, a volume evolving as $V \propto e^{\lambda\tau}$ leads to

$$\Theta(\varphi) \propto \varphi^{\frac{2(\Delta-1)+\Gamma}{(4-3\Gamma)\Delta}}, \quad (17)$$

that is, $\omega(\varphi) \propto \frac{1}{\varphi}$ and $\varphi \propto e^{\lambda\Delta\tau}$. To sum up, the only constraint imposed by assumption (14) is on the couple $(V(\tau), \omega(\varphi))$, through the relation:

$$\frac{\rho}{\rho_\varphi} = \frac{4(\Delta + 2)}{2(1 - \Delta) - \Gamma} V^{-2}, \quad (18)$$

that implies that, at any time, the ratio of the densities of the barotropic fluid and of the scalar field scales with the inverse of the square of the volume. This is a sufficient constraint to make the Kovalewski formalism tractable in scalar-tensor gravity. Replacing (15) into (13), and considering (14) after (9), the dynamical system associated with the homogeneous Universe in scalar-tensor theory is:

$$\begin{cases} E_c + E_p &= 2a^2 V^{2\Delta} + 2\varphi_o'^2 V^{2\Delta+2} \\ 2[2 - \Gamma] a^2 V^{2\Delta} &= A_1'' + n_1^2 \alpha_1^2 - (n_2 \alpha_2 - n_3 \alpha_3)^2 \\ 2[2 - \Gamma] a^2 V^{2\Delta} &= A_2'' + n_2^2 \alpha_2^2 - (n_3 \alpha_3 - n_1 \alpha_1)^2 \\ 2[2 - \Gamma] a^2 V^{2\Delta} &= A_3'' + n_3^2 \alpha_3^2 - (n_1 \alpha_1 - n_2 \alpha_2)^2 \end{cases} \quad (19)$$

It is important to note that whereas this system seems independent on Γ , it actually depends on it through the parameter Δ that fully characterizes our solution. In fact, for each coupling function, there exists a non-ambiguous link between Γ and Δ . For example in the case of a radiation fluid when $\Gamma = 4/3$, the LHS of the last equation in system (10) vanishes and one can find that $\varphi' \propto V^{-2}$ and then $\Delta = -2$.

A direct inspection of the first equation of system (19) shows that we can predict qualitatively the behavior of the case: $\Delta = 0$. Indeed, in this case, one of the two terms of $E_c + E_p$ reduces to a positive constant and the other one tends to 0 as V tends to 0. It is well known (e.g. [10]) that such a case breaks the Kasner cycle (e.g. [11]) for B_{VIII} and B_{IX} , and then suppresses the chaotic behaviour toward the $t = 0$ singularity for these models. The case $\Delta = -1$ seems to be similar, but the other term now diverges and then this simple analysis cannot be done.

1.4. Hamiltonian formalism

The quantity called $E_c = A'_1 A'_2 + A'_1 A'_3 + A'_3 A'_2$ is a quadratic form of $A_{i=1,2,3}$ derivatives. Then, one can diagonalize it using a linear change of variables:

$$\begin{cases} q_1 = (A_1 - A_2) / \sqrt{2} \\ q_2 = (A_1 + A_2 - 2A_3) / \sqrt{6} \\ q_3 = 2(A_1 + A_2 + A_3) / \sqrt{6} \end{cases} \quad (20)$$

Introducing the associated conformal time derivatives $p_{i=1,2,3} := q'_{i=1,2,3}$, the first equation of the system (19) becomes:

$$H := \frac{1}{2} (p_3^2 - p_1^2 - p_2^2) + e^{\frac{\sqrt{6}}{3} q_3} \xi(q_1, q_2) - 2a^2 e^{\frac{\sqrt{6}}{2} \Delta q_3} - 2\varphi_o'^2 e^{\frac{\sqrt{6}}{2} (\Delta+1) q_3} = 0 \quad (21)$$

the so-called potential ξ is defined by:

$$\begin{aligned} \xi(q_1, q_2) = & -n_1^2 e^{\left(\frac{\sqrt{6}}{3} q_2 + \sqrt{2} q_1\right)} - n_2^2 e^{\left(\frac{\sqrt{6}}{3} q_2 - \sqrt{2} q_1\right)} - n_3^2 e^{-\frac{2\sqrt{6}}{3} q_2} \\ & + 2n_1 n_2 e^{\frac{\sqrt{6}}{3} q_2} + 2n_1 n_3 e^{\left(\frac{\sqrt{2}}{2} q_1 - \frac{\sqrt{6}}{6} q_2\right)} + 2n_2 n_3 e^{-\left(\frac{\sqrt{2}}{2} q_1 + \frac{\sqrt{6}}{6} q_2\right)} \end{aligned} \quad (22)$$

In terms of (q_i, p_i) variables, the dynamical system (19) is quasi-Hamiltonian (e.g. [12] and later [13],[14],[15])

$$\begin{aligned} q'_{1,2} = \frac{dq_{1,2}}{d\tau} = -\frac{\partial H}{\partial p_{1,2}} \quad p'_{1,2} = \frac{dp_{1,2}}{d\tau} = -\frac{\partial H}{\partial q_{1,2}} \\ q'_3 = \frac{dq_3}{d\tau} = \frac{\partial H}{\partial p_3} \quad p'_3 = \frac{dp_3}{d\tau} = -\frac{\partial H}{\partial q_3} \end{aligned} \quad (23)$$

The minus that breaks the strict Hamiltonian symmetry comes from the minus signature of the quadratic form associated with E_c .

2. Integrability of homogeneous Universes

2.1. The case of Bianchi Universes

The following work has been initiated by Melnikov's team (see e.g. [16] and [17] and references therein). In the special case of B_{IX} , [18] consists in an application; a generalization in the context of the whole class A (i.e. $a = 0$) was tried by [19]. However, a lot of imprecisions in this last work need this new reformulation and extension to scalar-tensor theory.

2.1.1. Bianchi universes as generalized Toda systems Introducing the following vectors:

$$\begin{aligned} \mathbf{a}_1 &:= [0, \sqrt{6}/3, \sqrt{6}/3] & \mathbf{a}_2 &:= [\sqrt{2}/2, -\sqrt{6}/6, \sqrt{6}/3] & \mathbf{a}_3 &:= [-\sqrt{2}/2, -\sqrt{6}/6, \sqrt{6}/3] \\ \mathbf{a}_4 &:= [\sqrt{2}, \sqrt{6}/3, \sqrt{6}/3] & \mathbf{a}_5 &:= [-\sqrt{2}, \sqrt{6}/3, \sqrt{6}/3] & \mathbf{a}_6 &:= [0, -2\sqrt{6}/3, \sqrt{6}/3] \\ \mathbf{a}_7 &:= [0, 0, \sqrt{6}\Delta/2] & & & \mathbf{a}_8 &:= [0, 0, \sqrt{6}(\Delta+1)/2] \end{aligned} \quad (24)$$

the 3-forms:

$$\begin{aligned} \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^3 \quad (x, y) &:= +x_1 y_1 + x_2 y_2 + x_3 y_3 \\ \langle x, y \rangle &:= -x_1 y_1 - x_2 y_2 + x_3 y_3 \end{aligned} \quad (25)$$

and the constants:

$$\begin{aligned} k_1 &:= 2n_1n_2 & k_2 &:= 2n_1n_3 & k_3 &:= 2n_2n_3 \\ k_4 &:= -n_1^2 & k_5 &:= -n_2^2 & k_6 &:= -n_3^2 \\ k_7 &= -2a^2 & k_8 &= -2\varphi_o'^2 \end{aligned} \quad (26)$$

it is clear that the Hamiltonian (21) reads:

$$H = \frac{1}{2} \langle \mathbf{p}, \mathbf{p} \rangle + \sum_{i=1}^8 k_i e^{(\mathbf{a}_i, \mathbf{q})} \quad (27)$$

which is a classical form of a generalized Toda dynamical system.

Following Melnikov [16], we change $\{\mathbf{q}, \mathbf{p}\}$ variables to $\{\mathbf{u}, \mathbf{v}\}$ ones, such that:

$$\begin{cases} \mathbf{u} \in \mathbb{R}^8, u_{i=1,\dots,8} := \langle \mathbf{a}_i, \mathbf{p} \rangle \\ \mathbf{v} \in \mathbb{R}^8, v_{i=1,\dots,8} := \exp(\mathbf{a}_i, \mathbf{q}) \end{cases} \quad (28)$$

Through this change, the number of degrees of freedom jumps from 6 in $\{\mathbf{q}, \mathbf{p}\}$ to 16 in terms of $\{\mathbf{u}, \mathbf{v}\}$. Still writing ' for the derivatives with respect to the conformal time τ , the dynamical equations then read:

$$\forall i = 1, \dots, 8 \quad \begin{cases} v_i' = u_i v_i \\ u_i' = \sum_{j=1}^8 m_{ij} v_j \end{cases} \quad \text{with } m_{ij} := -k_j \langle \mathbf{a}_i, \mathbf{a}_j \rangle \quad (29)$$

This new formulation is now polynomial. Using the appendix notations, one can directly prove that the system (29) is autosimilar with any non vanishing index and weight $\mathbf{g}$ such that:

$$g_1 = \dots = g_8 = 1 \quad \text{and} \quad g_9 = \dots = g_{16} = 2 \quad (30)$$

$\mathbf{g}$ is unique provided that (A.3) is fulfilled. A particular autosimilar solution of (29) is then

$$[\tilde{\mathbf{u}} \ \tilde{\mathbf{v}}]^T = \mathbf{c} t^{-\mathbf{g}} = [\lambda_1 t^{-1}, \dots, \lambda_8 t^{-1}, \mu_1 t^{-2}, \dots, \mu_8 t^{-2}]^T \quad (31)$$

provided that the constant non vanishing vector $\mathbf{c} = [\lambda, \mu] = [\lambda_1, \dots, \lambda_8, \mu_1, \dots, \mu_8]$ is a solution of the algebraic system of equations:

$$\forall i = 1, \dots, 8 \quad \begin{cases} \sum_{j=1}^8 m_{ij} \mu_j = -\lambda_i \\ \lambda_i \mu_i = -2\mu_i \end{cases} \quad (32)$$

Hence, to any non vanishing solution of this last system corresponds a set of 16 Kovalewski exponents that allows to write the solution of the system (23) (cf Appendix A). A necessary condition for the system to be integrable is that all its Kovalewski exponents be rational. So, the rest of the paper will be devoted to the analysis of these exponents in order to study the integrability of different types of homogeneous and anisotropic Universes.

2.1.2. Solutions of the algebraic system In the more general case (that is scalar-tensor theory in presence of matter barotropic fluids), the algebraic system (32) makes use of the 8×8 matrix

$$M := \begin{bmatrix} 0 & -2n_1n_3 & -2n_3n_2 & 0 & 0 & 2n_3^2 & 2a^2\Delta & 2\varphi_o'^2\Delta_1 \\ -2n_1n_2 & 0 & -2n_3n_2 & 0 & 2n_2^2 & 0 & 2a^2\Delta & 2\varphi_o'^2\Delta_1 \\ -2n_1n_2 & -2n_1n_3 & 0 & 2n_1^2 & 0 & 0 & 2a^2\Delta & 2\varphi_o'^2\Delta_1 \\ 0 & 0 & -4n_3n_2 & -2n_1^2 & 2n_2^2 & 2n_3^2 & 2a^2\Delta & 2\varphi_o'^2\Delta_1 \\ 0 & -4n_1n_3 & 0 & 2n_1^2 & -2n_2^2 & 2n_3^2 & 2a^2\Delta & 2\varphi_o'^2\Delta_1 \\ -4n_1n_2 & -4n_1n_3 & 0 & 2n_1^2 & 2n_2^2 & -2n_3^2 & 2a^2\Delta & 2\varphi_o'^2\Delta_1 \\ -2n_1n_2\Delta & -2n_1n_3\Delta & -2n_3n_2\Delta & n_1^2\Delta & n_2^2\Delta & n_3^2\Delta & 3a^2\Delta^2 & 3\varphi_o'^2\Delta_1\Delta \\ -2n_1n_2\Delta_1 & -2n_1n_3\Delta_1 & -2n_3n_2\Delta_1 & n_1^2\Delta_1 & n_2^2\Delta_1 & n_3^2\Delta_1 & 3a^2\Delta\Delta_1 & 3\varphi_o'^2\Delta_1^2 \end{bmatrix} \quad (33)$$

where $\Delta_1 := \Delta + 1$. One can straightforwardly check that M is of rank 3.

The vector 0 is always a solution of the system (32), but it is not relevant for the quest of Kovalewski exponents that needs non trivial solutions. A systematic study of the solutions of the system (32) is possible : for $p = 6, 7$ or 8 let $E_p = \{1, 2, 3, \dots, p\}$, we consider the minor determinants:

$$\forall (i, j, k) \in E_p \times E_p \times E_p \quad \zeta_i = m_{ii}, \zeta_{i,j} = \begin{vmatrix} m_{ii} & m_{ij} \\ m_{ji} & m_{jj} \end{vmatrix} \\ \zeta_{i,j,k} = \begin{vmatrix} m_{ii} & m_{ij} & m_{ik} \\ m_{ji} & m_{jj} & m_{jk} \\ m_{ki} & m_{kj} & m_{kk} \end{vmatrix} \quad (34)$$

and the determinants:

$$d_{2i} = \begin{vmatrix} 2 & m_{ij} \\ 2 & m_{jj} \end{vmatrix}, \quad d_{2j} = \begin{vmatrix} m_{ii} & 2 \\ m_{ji} & 2 \end{vmatrix} \quad (35)$$

$$d_{3i} = \begin{vmatrix} 2 & m_{ij} & m_{ik} \\ 2 & m_{jj} & m_{jk} \\ 2 & m_{kj} & m_{kk} \end{vmatrix}, \quad d_{3j} = \begin{vmatrix} m_{ii} & 2 & m_{ik} \\ m_{ji} & 2 & m_{jk} \\ m_{ki} & 2 & m_{kk} \end{vmatrix}, \quad d_{3k} = \begin{vmatrix} m_{ii} & m_{ij} & 2 \\ m_{ji} & m_{jj} & 2 \\ m_{ki} & m_{kj} & 2 \end{vmatrix} \quad (36)$$

Solutions of System (32) are then classified into three classes:

- (i) Type 1 solutions (T1): $\exists! i \in E_p$ such that $\mu_i \neq 0$ and $\forall j \in E_p \setminus \{i\}, \mu_j = 0$: then $\lambda_i = -2$ and
 - if $\zeta_i = 0$: there is no solution.
 - if $\zeta_i \neq 0$: $\mu_i = 2/\zeta_i$ and $\forall j \in E_p \setminus \{i\}, \lambda_j = -2m_{ji}/\zeta_i$
- (ii) Type 2 solutions (T2): $\exists! (i, j) \in E_p \times (E_p \setminus \{i\})$ such that $\{\mu_i, \mu_j\} \neq \{0, 0\}$ and $\forall k \in E_p \setminus \{i, j\}, \mu_k = 0$: then $\lambda_i = \lambda_j = -2$ and
 - if $\zeta_{i,j} = 0$: there is no solution.
 - if $\zeta_{i,j} \neq 0$: $\mu_i = d_{2i}/\zeta_{ij}$ and $\mu_j = d_{2j}/\zeta_{i,j}$ moreover $\forall k \in E_p \setminus \{i, j\}, \lambda_k = (m_{ki}d_{2i} + m_{kj}d_{2j})/\zeta_{i,j}$
- (iii) Type 3 solutions (T3): $\exists! (i, j, k) \in E_p \times (E_p \setminus \{i\}) \times (E_p \setminus \{i, j\})$ such that $\{\mu_i, \mu_j, \mu_k\} \neq \{0, 0, 0\}$ and $\forall l \in E_p \setminus \{i, j, k\}, \mu_l = 0$: then $\lambda_i = \lambda_j = \lambda_k = -2$ and

- if $\zeta_{i,j,k} = 0$: there is no solution
- if $\zeta_{i,j,k} \neq 0$: $\mu_i = d_{3i}/\zeta_{i,j,k}$, $\mu_j = d_{3j}/\zeta_{i,j,k}$ and $\mu_k = d_{3k}/\zeta_{i,j,k}$ moreover $\forall l \in E_p \setminus \{i, j, k\}$, $\lambda_l = (m_{li}d_{3i} + m_{lj}d_{3j} + m_{lk}d_{3k})/\zeta_{i,j,k}$

2.1.3. Kovalewski exponents for Bianchi Universes In what follows, we will examine the Kovalewski exponents for Bianchi Universes in 4 different cases that are included in the formalism presented above: scalar-tensor gravity with or without matter fluids, and General Relativity with or without matter fluids. We proceed in three steps.

- (i) Select a Universe. It corresponds to choosing a set of $n_{i=1,2,3}$ in table 1.1. This step determines the theory of gravitation and the matter content of interest, and this choice is made by considering different forms for the matrix M defined in (i):

- Barotropic matter filled Universe in scalar-tensor theory of gravitation: $p = 8$. This is the most general case that was presented in detail in the preceding subsection.
- Empty Universe in scalar-tensor theory of gravitation: $p = 7$. The associated matrix is M but without the 7th line and the 7th column:

$$M_{STV} := \begin{bmatrix} 0 & -2n_1n_3 & -2n_3n_2 & 0 & 0 & 2n_3^2 & 2\varphi_o'^2\Delta_1 \\ -2n_1n_2 & 0 & -2n_3n_2 & 0 & 2n_2^2 & 0 & 2\varphi_o'^2\Delta_1 \\ -2n_1n_2 & -2n_1n_3 & 0 & 2n_1^2 & 0 & 0 & 2\varphi_o'^2\Delta_1 \\ 0 & 0 & -4n_3n_2 & -2n_1^2 & 2n_2^2 & 2n_3^2 & 2\varphi_o'^2\Delta_1 \\ 0 & -4n_1n_3 & 0 & 2n_1^2 & -2n_2^2 & 2n_3^2 & 2\varphi_o'^2\Delta_1 \\ -4n_1n_2 & -4n_1n_3 & 0 & 2n_1^2 & 2n_2^2 & -2n_3^2 & 2\varphi_o'^2\Delta_1 \\ -2n_1n_2\Delta_1 & -2n_1n_3\Delta_1 & -2n_3n_2\Delta_1 & n_1^2\Delta_1 & n_2^2\Delta_1 & n_3^2\Delta_1 & 3\varphi_o'^2\Delta_1^2 \end{bmatrix}$$

- Barotropic matter filled Universe in Einstein General Relativity: $p = 7$. The associated matrix is M but with the 8th line and 8th column removed, and the 7th line and 7th column adapted. Noting γ for $2 - \Gamma$ we have:

$$M_{GRM} := \begin{bmatrix} 0 & -2n_1n_3 & -2n_3n_2 & 0 & 0 & 2n_3^2 & 2\rho_o\chi\gamma \\ -2n_1n_2 & 0 & -2n_3n_2 & 0 & 2n_2^2 & 0 & 2\rho_o\chi\gamma \\ -2n_1n_2 & -2n_1n_3 & 0 & 2n_1^2 & 0 & 0 & 2\rho_o\chi\gamma \\ 0 & 0 & -4n_3n_2 & -2n_1^2 & 2n_2^2 & 2n_3^2 & 2\rho_o\chi\gamma \\ 0 & -4n_1n_3 & 0 & 2n_1^2 & -2n_2^2 & 2n_3^2 & 2\rho_o\chi\gamma \\ -4n_1n_2 & -4n_1n_3 & 0 & 2n_1^2 & 2n_2^2 & -2n_3^2 & 2\rho_o\chi\gamma \\ -n_1n_2\gamma & -n_1n_3\gamma & -n_3n_2\gamma & n_1^2\gamma/2 & n_2^2\gamma/2 & n_3^2\gamma/2 & 3\rho_o\chi\gamma^2/2 \end{bmatrix}$$

- Empty Universe in Einstein General Relativity: $p = 6$. The associated matrix is M but with the 7th and 8th lines and the 7th and 8th columns removed:

$$M_{GRV} := \begin{bmatrix} 0 & -2n_1n_3 & -2n_3n_2 & 0 & 0 & 2n_3^2 \\ -2n_1n_2 & 0 & -2n_3n_2 & 0 & 2n_2^2 & 0 \\ -2n_1n_2 & -2n_1n_3 & 0 & 2n_1^2 & 0 & 0 \\ 0 & 0 & -4n_3n_2 & -2n_1^2 & 2n_2^2 & 2n_3^2 \\ 0 & -4n_1n_3 & 0 & 2n_1^2 & -2n_2^2 & 2n_3^2 \\ -4n_1n_2 & -4n_1n_3 & 0 & 2n_1^2 & 2n_2^2 & -2n_3^2 \end{bmatrix}$$

We then have to apply the following algorithm to the appropriate matrix.

- (ii) Determine all the non vanishing minor determinants ζ which could be extracted from the considered matrix.

- (iii) For each ζ , compute the associated set of $2p$ Kovalewski exponents. In our polynomial case, as indicated in appendix A this set is the set of eigenvalues of the matrix:

$$K = \begin{bmatrix} 1 & 0 & \cdots & \cdots & 0 & m_{11} & \cdots & \cdots & \cdots & m_{p1} \\ 0 & \ddots & \ddots & & \vdots & \vdots & & & & \vdots \\ \vdots & \ddots & I_p & \ddots & \vdots & \vdots & & M & & \vdots \\ \vdots & & \ddots & \ddots & 0 & \vdots & & & & \vdots \\ 0 & \cdots & \cdots & 0 & 1 & m_{1p} & \cdots & \cdots & \cdots & m_{pp} \\ \mu_1 & 0 & \cdots & \cdots & 0 & \lambda_1 + 2 & 0 & \cdots & \cdots & 0 \\ 0 & \ddots & \ddots & & \vdots & 0 & \ddots & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 & \vdots & & \ddots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & \mu_p & 0 & \cdots & \cdots & 0 & \lambda_p + 2 \end{bmatrix} \quad (37)$$

The particular form of the matrix K and the weak rank of the matrix M allows an explicit calculation of the eigenvalues in all the cases $p = 6, 7, 8$ and for all the Bianchi Universes. There exists several hundreds of non vanishing minor determinants in all the cases considered. They correspond to 92 distinct sets of Kovalewski exponents with length $2p = 12, 14$ or 16 . All the details are presented in appendix B. In all cases, calculations are explicit, and in order to organize it we use formal calculus computational tools.

2.1.4. Exponents analysis and conclusions The full set of Kovalewski exponents is presented in appendix B. The analysis of the sets of exponents results in four classes for Bianchi models : Class I contains uniquely B_I models, Class II contains B_{II} and B_{IV} , Class III is composed by B_{III} , $B_{VI_{o,a}}$ and $B_{VII_{o,a}}$, and finally Class IV contains B_{VIII} and the famous B_{IX} . Each class is characterized by the same set of Kovalewski exponents, therefore corresponding models have the same characteristic dynamics. Let us review for each class the algebraic properties of Kovalewski exponents :

- Class I — For all the cases we studied, all the Kovalewski exponents belonging to this class, are rational provided that the barotropic index Γ and/or the scalar-tensor parameter Δ are rational. This restriction corresponds to physical cases ‡ .
- Class II — Kovalewski exponents of models belonging to this class fall into four subcases :
 - Empty Universe in General Relativity (EUGR): exponents belonging to this class are integers.
 - Barotropic Matter filled Universe in General Relativity (BMUGR): due to $A_\pm^*$, all exponents are rational iff $\Gamma \in \mathbb{Q} \cap [0, \Gamma_0]$ with $\Gamma_0 = (11 - \sqrt{73})/3 \approx 0.81$.

‡ As a matter of fact, physical power laws or barotropic index must be rational in order to be full of physical meaning

- Empty Universe in Scalar-Tensor Theory (EUSTT): due to $A_{\pm}$, all exponents are rational iff $\Delta \in \mathbb{Q} \cap [\Delta_0, 1]$ with $\Delta_0 = (-11 + \sqrt{73})/6 \approx -0.40$.
- Barotropic Matter filled Universe in Scalar-Tensor Theory (BMUSTT): Due to $A_{\pm}$ and $D_{\pm}$, all exponents are rational iff $\Delta \in \mathbb{Q} \cap [\Delta_1, 1]$ with $\Delta_1 = (-5 + \sqrt{73})/6 \approx 0.59$.
- Class III – As previously, four classes of sets of Kovalewski exponents can be identified
 - EUGR : exponents belonging to this class are integers.
 - BMUGR : due to $A_{\pm}^*$, $B_{\pm}^*$ and $C_{\pm}^*$, all exponents are rational iff $\Gamma \in \mathbb{Q} \cap [0, \Gamma_0]$. This case is then equivalent to the corresponding case of class II models. Let us remark the special value $\Gamma = 2/3$ for which $(A_+^*, A_-^*, B_+^*, B_-^*, C_+^*, C_-^*) = (1, 0, 1, 0, 1, 0)$.
 - EUSTT : due to $A_{\pm}, B_{\pm}$ and $a_{\pm}$ all exponents are rational iff $\Delta \in \mathbb{Q} \cap [\Delta_0, 1]$. This case is then equivalent to the corresponding case of class II models. Let us remark the special value $\Delta = -1/3$ for which $(A_+, A_-, B_+, B_-, a_+, a_-) = (1, 0, 1, 0, 1, 0)$.
 - BMUSTT : due to $A_{\pm}, B_{\pm}, C_{\pm}, D_{\pm}, a_{\pm}$ and $b_{\pm}$, all exponents are rational iff $\Delta \in \mathbb{Q} \cap [\Delta_2, 1]$ with $\Delta_2 = 16/25 = 0.64$.
- Class IV – all models of this class contain at least two conjugated complex Kovalewski exponents.

From such an analysis, taking into account Yoshida's theorems (see [21, 22]) whose context is detailed in appendix A, we can conclude that:

- Empty Universes whose metrics correspond to classes I, II or III defined below are generically associated to algebraically integrable dynamics.
- Class IV Universes are always associated to non algebraically integrable dynamics. This result holds for General Relativity and/or scalar-tensor theory we are interested in, the Universe being empty and/or filled of barotropic matter.
- BMUGR of class II and III are associated to non integrable dynamics if the barotropic index ranges in the interval $[\Gamma_0, 2]$ with $\Gamma_0 = (11 - \sqrt{73})/3 \approx 0.81$.
- EUSTT of class II and III are associated to non integrable dynamics if the power law of the scalar-tensor modelization (14) ranges in the interval $[-2, \Delta_0]$ with $\Delta_0 = (-11 + \sqrt{73})/6 \approx -0.40$.
- BMUSTT of class II are associated to non integrable dynamics if the power law of the scalar tensor modelization (see 14) ranges in the interval $[-2, \Delta_1]$ with $\Delta_1 = (-5 + \sqrt{73})/6 \approx -0.40$.
- BMUSTT of class III are associated to non integrable dynamics if the power law of the scalar tensor modelization (14) ranges in the interval $[-2, \Delta_2]$ with $\Delta_2 = 16/25 = 0.64$.

The integrability of homogeneous Universes depends on multiple factors.

It is well known, since the pioneering works by Elskens and Henneaux [23], that the number of dimensions D of the spatial sections of the Universe is one of this factors. It influences the form of the potential $\xi(q)$ in relation (22) or/and increases the number of components of the vectors $\{\mathbf{a}_i\}$ in (24). The net result in the simplified context of generalized Kasner metric is that chaos disappears – and then system becomes always integrable, in a classical context – when $D \geq 11$.

Modification of the gravitation theory by introducing a scalar-tensor component seems here to have a different contribution. As a matter of fact, in the case we consider, the potential $\xi(q)$ is not modified, but a volume dependent Toda potential is added in the Hamiltonian. In a general way, this produces a slowing down for the cushions' velocity in the billiard representation (see [15] and references therein). One can then reasonably suppose that this feature does not change drastically the dynamical properties of homogeneous Universes inherited from General Relativity. The integrability analysis based on Kovalewski exponents we have done confirms such a conjecture.

Appendix A : Integrability of autosimilar differential systems

Let $\mathbf{f}$ be a continuous function from $\mathbb{R}^n$ to $\mathbb{R}^n$ and $\mathbf{x} = [x_1(t), \dots, x_n(t)]^T \in \mathbb{R}^n$ such that

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}) \quad (\text{A.1})$$

If there exists a positive real λ and a rational vector $\mathbf{g} := [g_1, \dots, g_n]^T$ such that the transformation

$$\begin{aligned} t &\mapsto t/\lambda \\ \forall i = 1, \dots, n \quad x_i &\mapsto \lambda^{g_i} x_i \end{aligned} \quad (\text{A.2})$$

leaves the system (A.1) invariant, this system is called autosimilar with index λ and weight $\mathbf{g}$.

When it exists $\mathbf{g}$, it is unique if

$$A(\mathbf{x}) \in M_n(\mathbb{R}) \quad , \quad A_{ij} = x_j \frac{\partial f_i(\mathbf{x})}{\partial x_j} - \delta_{ij} f_i(\mathbf{x}) \quad (\text{A.3})$$

is invertible for almost all $\mathbf{x} \in \mathbb{R}^n$.

Autosimilar systems always admit at least one autosimilar particular solution

$$\tilde{\mathbf{x}}_{as} = [c_1 (t - t_o)^{-g_1}, \dots, c_n (t - t_o)^{-g_n}]^T \quad (\text{A.4})$$

where $\mathbf{c} := [c_1, \dots, c_n]^T$ is solution of the algebraic equation:

$$\begin{cases} f_1(\mathbf{c}) = -g_1 c_1 \\ \vdots \\ f_n(\mathbf{c}) = -g_n c_n \end{cases} \quad (\text{A.5})$$

Linearization of the system (A.1) around the solution $\tilde{\mathbf{x}}_{as}$ gives

$$\frac{d\mathbf{z}}{dt} = D\mathbf{f}(\mathbf{x})(\tilde{\mathbf{x}}_{as}) \mathbf{z} \quad (\text{A.6})$$

where $D\mathbf{f}(\mathbf{x})$ ($\tilde{\mathbf{x}}_{as}$) is the Jacobian matrix of $\mathbf{f}(\mathbf{x})$ evaluated at $\mathbf{x} = \tilde{\mathbf{x}}_{as}$. A theorem by Fuchs (see [20] §5.6 and 5.7) then shows that the general solution of (A.6) is autosimilar and reads:

$$\mathbf{z} = [k_1 (t - t_o)^{\rho_1 - g_1}, \dots, k_n (t - t_o)^{\rho_n - g_n}]^T \quad (\text{A.7})$$

The quantities $(\rho_1, \rho_2, \dots, \rho_n)$ are called Kovalewski exponents. More practically, one can compute them because there are also eigenvalues of the matrix:

$$K := D\mathbf{f}(\mathbf{x})(\mathbf{c}) + \text{diag}(\mathbf{g}). \quad (\text{A.8})$$

As shown by Comte (see [20] §5.6 and 5.7), they are of great importance for the study of integrability of the original non linear system (A.1). As a matter of fact each component $x_{i=1,\dots,n}$ of a solution x of this system can be written as:

$$x_i(t) = \sum_{k=0}^{+\infty} \varepsilon^k x_i^{(k)}(t) \quad \text{with} \quad \begin{cases} x_i^{(0)} = c_i (t - t_o)^{g_i} \\ x_i^{(1)} = k_i (t - t_o)^{\rho_i - g_i} \\ \exists 1 \leq p, q \leq n, \quad x_i^{(2)} = c_p k_q (t - t_o)^{g_p + \rho_q - g_q} \\ \text{etc} \dots \end{cases} \quad (\text{A.9})$$

hence

$$x_i(t) \propto (t - t_o)^{g_i} S[(t - t_o)^{\rho_1}, \dots, (t - t_o)^{\rho_n}] \quad (\text{A.10})$$

where $S[]$ is a multiple series. This allows to understand a theorem by Yoshida (see [21, 22]) : a necessary condition for a differential system to be integrable is that all its Kovalewski exponents are rational. If there exists at least one exponent irrational or complex the corresponding differential system is not algebraically integrable.

Appendix B : Details of Kovalewski exponents

When only non exotic barotropic matter ($0 \leq \Gamma < 2$) is considered, we have noted $\gamma = (3\Gamma - 2)/(\Gamma - 2)$. For any $K = 3, 5, 7$ and 15 we note $2K^{\pm i} = 1 \pm i\sqrt{K}$ and

$$\begin{aligned} 2A_{\pm}^* &= 1 \pm \sqrt{\frac{16-22\Gamma+3\Gamma^2}{4-2\Gamma}} & 2B_{\pm}^* &= 1 \pm \sqrt{5-6\Gamma} \\ 2C_{\pm}^* &= 1 \pm \sqrt{\frac{25\Gamma-18}{\Gamma-2}} & 2D_{\pm}^* &= 1 \pm \sqrt{\frac{18-41\Gamma+24\Gamma^2}{2-\Gamma}} \end{aligned}$$

$$E_{\gamma} := \{-1, 2, 1(\times 6), \frac{2\gamma}{3}(\times 6)\} \quad E_o := \{-1, 4(\times 3), 2(\times 3), 1(\times 5)\}$$

Moreover

$$E_{\delta} = \{-1, 1(\times 6), 2, 2\delta_1/3(\times 6)\}$$

$$\delta_1 = (3\Delta + 1)(\Delta + 1)^{-1} \quad \text{and} \quad \delta_2 = (3\Delta - 2)\Delta^{-1}$$

$$2A_{\pm=1} = 1 \pm \sqrt{\frac{3(\Delta - \Delta_A^+)(\Delta - \Delta_A^-)}{|\Delta + 1|}} \quad 6\Delta_A^{\pm} = -11 \pm \sqrt{73}$$

$$2D_{\pm} = 1 \pm \sqrt{\frac{3(\Delta - \Delta_D^+)(\Delta - \Delta_D^-)}{|\Delta|}} \quad 6\Delta_D^{\pm} = -5 \pm \sqrt{73}$$

$$2B_{\pm} = 1 \pm \sqrt{\frac{25\Delta + 9}{|\Delta + 1|}}$$

$$2C_{+} = 1 \pm \sqrt{\frac{25\Delta - 16}{|\Delta|}}$$

$$2E_{\pm} = 1 \pm \sqrt{\frac{48\Delta^2 + 41\Delta + 9}{|\Delta + 1|}} \quad 48\Delta^2 + 41\Delta + 9 > 0$$

$$2F_{\pm} = 1 \pm \sqrt{\frac{48\Delta^2 - 55\Delta + 16}{|\Delta|}} \quad 48\Delta^2 - 55\Delta + 16 > 0$$

$$2a_{\pm} = 1 \pm \sqrt{5 + 12\Delta} \quad 2b_{\pm} = 1 \pm \sqrt{-7 + 12\Delta}$$

Using the notations defined above we have computed all the Kowalewski exponents for all Bianchi Universes, in the case of empty or barotropic filled Universes and for General Relativity (GR) and Scalar-Tensor (ST) theory of gravitation:

B_I		
RG	Empty (N=6)	Barotropic Matter (N=7)
T1	$\emptyset$	E_{γ}
T2	$\emptyset$	$\emptyset$
T3	$\emptyset$	$\emptyset$
ST	Empty (N=7)	Barotropic Matter (N=8)
T1	E_{δ}	$-1, 1(\times 7), 2, 2\delta_2/3(\times 6), -2/\Delta$ $-1, 1(\times 7), 2, 2(\Delta + 1)^{-1}, 2\delta_1/3(\times 6)$
T2	$\emptyset$	$\emptyset$
T3	$\emptyset$	$\emptyset$

B_{II} and B_{IV}		
GR	Empty (N=6)	Barotropic Matter (N=7)
T1	E_o	E_{γ} $-1, 1(\times 6), 2(\times 3), 4(\times 3), 3 - \Gamma/2$
T2	$\emptyset$	$-1, 1(\times 5), 2, \gamma/2(\times 2), \gamma(\times 3), A_{\pm}^*$
T3	$\emptyset$	$\emptyset$
ST	Empty (N=7)	Barotropic Matter (N=8)
T1	$-1, 1(\times 6), 2(\times 3), 4(\times 3), \Delta + 3$ E_{δ}	$-1, 1(\times 7), 2, 2(\Delta + 1)^{-1}, 2\delta_1/3(\times 6)$ $-1, 1(\times 7), 2, 2\delta_2/3(\times 6), -2/\Delta$ $-1, 1(\times 7), 2(\times 3), 4(\times 3), \Delta + 3, \Delta + 2$
T2	$-1, 1(\times 5), 2, \delta_1/2(\times 2), \delta_1(\times 3), A_{\pm}$	$-1, 1(\times 6), 2, \delta_2/2(\times 2), \delta_2(\times 3), -2/\Delta, D_{\pm}$ $-1, 1(\times 6), 2, \delta_1/2(\times 2), \delta_1(\times 3), 2(\Delta + 1)^{-1}, A_{\pm}$
T3	$\emptyset$	$\emptyset$

$B_{III}, B_{VI_{0,a}}, B_{VII_{0,a}}$		
GR	Empty (N=6)	Barotropic Matter(N=7)
T1	E_o	E_γ $-1, 1(\times 6), 2(\times 3), 4(\times 3), 3 - \Gamma/2$
T2	$\emptyset$	$-1, 1(\times 5), 2, \gamma/2(\times 2), \gamma(\times 3), A_\pm^*$ $-1, 0(\times 2), 1(\times 5), 2, 2\gamma, \gamma(\times 2), B_\pm^*$
T3	$\emptyset$	$-1, 0, 1(\times 4), 2, 2\gamma, \gamma(\times 2), B_\pm^*, C_\pm^*$ $-1, 0(\times 2), 1(\times 5), 2, 2\gamma, \gamma(\times 2), B_\pm^*$
ST	Empty (N=7)	Barotropic Matter (N=8)
T1	$-1, 1(\times 6), 2(\times 3), 4(\times 3), \Delta + 3$ E_δ	$-1, 1(\times 7), 2, 2\delta_1/3(\times 6), 2(\Delta + 1)^{-1}$ $-1, 1(\times 7), 2, 2\delta_2/3(\times 6), -2\Delta^{-1}$ $-1, 1(\times 7), 2(\times 3), 4(\times 3), \Delta + 3, \Delta + 2$
T2	$-1, 1(\times 5), 2, \delta_1/2(\times 2), \delta_1(\times 3), A_\pm$ $-1, 0(\times 2), 1(\times 5), 2, 2\delta_1, \delta_1(\times 2), a_\pm$	$-1, 0(\times 2), 1(\times 6), 2, 2\delta_1, \delta_1(\times 2), 2(\Delta + 1)^{-1}, a_\pm$ $-1, 1(\times 6), 2, \delta_1/2(\times 2), \delta_1(\times 3), 2(\Delta + 1)^{-1}, A_\pm$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_2, \delta_2(\times 2), -2\Delta^{-1}, b_\pm$ $-1, 1(\times 6), 2, \delta_2/2(\times 2), \delta_2(\times 3), -2\Delta^{-1}, D_\pm$
T3	$-1, 0, 1(\times 4), 2, 2\delta_1, \delta_1(\times 2), a_\pm, B_\pm$ $-1, 0(\times 2), 1(\times 5), 2, 2\delta_1, \delta_1(\times 2), a_\pm$	$-1, 0, 1(\times 5), 2, 2\delta_1, \delta_1(\times 2), 2(\Delta + 1)^{-1}, a_\pm, B_\pm$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_1, \delta_1(\times 2), 2(\Delta + 1)^{-1}, a_\pm$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_2, \delta_2(\times 2), -2\Delta^{-1}, b_\pm$ $-1, 0, 1(\times 5), 2, \delta_2(\times 2), 2\delta_2, -2\Delta^{-1}, b_\pm, C_\pm$

B_{VIII} and B_{IX}		
GR	Empty (N=6)	Barotropic Matter (N=7)
T1	E_o	E_γ $-1, 1(\times 6), 2(\times 3), 4(\times 3), 3 - \Gamma/2$
T2	$-1, -2(\times 3), 7^{\pm i}, 2(\times 2), 1(\times 4)$ $-1, 2, 15^{\pm i}, 1(\times 4), 0(\times 4)$	$-1, 1(\times 5), 2, \gamma/2(\times 2), \gamma(\times 3), A_\pm^*$ $-1, 0(\times 2), 1(\times 5), 2, 2\gamma, \gamma(\times 2), B_\pm^*$ $-2(\times 3), -1, 1(\times 5), 2(\times 2), 7^{\pm i}, 2\Gamma - 2$ $-1, 0(\times 4), 1(\times 5), 2, 15^{\pm i}, -1 + 3\Gamma/2$
T3	$-1, 2, 15^{\pm i}(\times 2), 1(\times 3), 0(\times 3)$ $-1, 2, 3^{\pm i}(\times 2), 1(\times 3), 0(\times 3)$ $-1, 2, 15^{\pm i}, 1(\times 4), 0(\times 4)$ $-1(\times 2), 2(\times 2), 7^{\pm i}, 1(\times 3), 0(\times 3)$	$-1, 0(\times 3), 1(\times 4), 2, 3^{\pm i}(\times 2), -1 + 3\Gamma/2$ $-1, 1(\times 4), 2, -2\gamma, 2\gamma(\times 3), C_\pm^*, D_\pm^*$ $-1, 0(\times 3), 1(\times 4), 2, 15^{\pm i}(\times 2), -1 + 3\Gamma/2$ $-1, 0(\times 4), 1(\times 5), 2, 15^{\pm i}, -1 + 3\Gamma/2$ $-1, 0, 1(\times 4), 2, 2\gamma, \gamma(\times 2), B_\pm^*, C_\pm^*$ $-1(\times 2), 0(\times 3), 1(\times 4), 2(\times 2), 7^{\pm i}, -1 + 3\Gamma/2$ $-1, 0(\times 2), 1(\times 5), 2, 2\gamma, \gamma(\times 2), B_\pm^*$
ST	Empty (N=7)	Barotropic Matter (N=8)
T1	E_δ $-1, 1(\times 6), 2(\times 3), 4(\times 3), \Delta + 3$	$-1, 1(\times 7), 2, 2\delta_2/3(\times 6), -2\Delta^{-1}$ $-1, 1(\times 7), 2(\times 3), 4(\times 3), \Delta + 3, \Delta + 2$ $-1, 1(\times 7), 2, 2\delta_1/3(\times 6), 2(\Delta + 1)^{-1}$
T2	$-1, 0(\times 2), 1(\times 5), 2, \delta_1(\times 2), 2\delta_1, a_\pm$ $-2(\times 3), -1, 1(\times 5), 2(\times 2), 7^{\pm i}, -4\Delta - 2$ $-1, 0(\times 4), 1(\times 5), 2, 15^{\pm i}, -3\Delta - 1$ $-1, 1(\times 5), 2, \delta_1/2(\times 2), \delta_1(\times 3), A_\pm$	$-1, 1(\times 6), 2, D_\pm, \delta_2/2(\times 2), \delta_2(\times 3), -2\Delta^{-1}$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_2, \delta_2(\times 2), -2\Delta^{-1}, b_\pm$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_1, \delta_1(\times 2), 2(\Delta + 1)^{-1}, a_\pm$ $-1, 1(\times 6), 2, \delta_1/2(\times 2), \delta_1(\times 3), 2(\Delta + 1)^{-1}, A_\pm$ $-2(\times 3), -1, 1(\times 6), 2(\times 2), 7^{\pm i}, -4\Delta - 2, -4\Delta + 2$ $-1, 0(\times 4), 1(\times 6), 2, 15^{\pm i}, -3\Delta - 1, -3\Delta + 2$
T3	$-1, 0(\times 2), 1(\times 5), 2, 2\delta_1, \delta_1(\times 2), a_\pm$ $-1, 0(\times 4), 1(\times 5), 2, 15^{\pm i}, -3\Delta - 1$ $-1, 0(\times 3), 1(\times 4), 2, 15^{\pm i}(\times 2), -3\Delta - 1$ $-1, 0(\times 3), 1(\times 4), 2, 3^{\pm i}(\times 2), -3\Delta - 1$ $-1, 0, 1(\times 4), 2, 2\delta_1, \delta_1(\times 2), a_\pm, B_\pm$ $-1(\times 2), 0(\times 3), 1(\times 4), 2(\times 2), 7^{\pm i}, -3\Delta - 1$ $-1, 1(\times 4), 2, -2\delta_1, 2\delta_1(\times 3), B_\pm, E_\pm$	$-1, 1(\times 5), 2, -2\delta_1, 2\delta_1(\times 3), 2(\Delta + 1)^{-1}, B_\pm, E_\pm$ $-1, 1(\times 5), 2, -2\delta_2, 2\delta_2(\times 3), -2\Delta^{-1}, C_\pm, F_\pm$ $-1, 0(\times 3), 1(\times 5), 2, 3^{\pm i}(\times 2), -3\Delta - 1, -3\Delta + 2$ $-1, 0(\times 3), 1(\times 5), 2, 15^{\pm i}(\times 2), -3\Delta - 1, -3\Delta + 2$ $-1(\times 2), 0(\times 3), 1(\times 5), 2(\times 2), 7^{\pm i}, -3\Delta - 1, -3\Delta + 2$ $-1, 0, 1(\times 5), 2, 2\delta_1, \delta_1(\times 2), 2(\Delta + 1)^{-1}, a_\pm, B_\pm$ $-1, 0, 1(\times 5), 2, 2\delta_2, \delta_2(\times 2), -2\Delta^{-1}, b_\pm, C_\pm$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_2, \delta_2(\times 2), -2\Delta^{-1}, b_\pm$ $-1, 0(\times 2), 1(\times 6), 2, 2\delta_1, \delta_1(\times 2), 2(\Delta + 1)^{-1}, a_\pm$ $-1, 0(\times 4), 1(\times 6), 2, 15^{\pm i}, -3\Delta - 1, -3\Delta + 2$

References

- [1] L. Bianchi, Sugli spazi a tre dimensioni che ammettono un gruppo continuo di movimenti [On three-dimensional spaces which admit a continuous group of motions], Memorie di Matematica e di Fisica della Societa Italiana delle Scienze, Serie Terza, vol. 11, p.267-352, 1898
- [2] L. Bianchi, Lezioni sulla teoria dei gruppi continui finite di trasformazioni [Lectures on the theory of finite continuous transformation groups (1902-1903)], Pisa, 1918, pp.550-557. English

- translation: General Relativity and Gravitation, 33, 2157-2170, 2001, and General Relativity and Gravitation, 33, 2171-2253, 2001
- [3] C. Brans and R. Dicke, Mach's Principle and a Relativistic Theory of Gravitation, *Phys. Rev.*, 124, 925, 1961
 - [4] P. Bergmann, Comments on the scalar tensor theory, *Int J. Theor. Phys.*, 1, 25, 1968
 - [5] K. Nordvedt, Post-Newtonian Metric for a General class of Scalar-Tensor Gravitational Theories and Observational Consequences, *Astrophys. J.*, 161, 1059, 1970
 - [6] R. Wagoner, Scalar tensor theory and gravitational waves, *Phys. Rev. D*, 1, 3209, 1970
 - [7] T. Damour and G. Esposito-Farèse, Tensor-multi-scalar theories of gravitation, *Class. Quant. Grav.*, 9, 2093-2176, 1992
 - [8] M. MacCallum, Anisotropic and Inhomogeneous Relativistic Cosmologies, in [9], p. 533-580, 1979
 - [9] S. Hawking and W. Israel, General relativity : an Einstein Centenary Survey, Cambridge University Press, 1979
 - [10] V. A. Belinskii and L.M. Khalatnikov, Effect of scalar and vector fields on the nature of the cosmological singularity, *Sov. Phys. JETP*, 36, 591, 1973
 - [11] V. A. Belinskii, L.M. Khalatnikov and E.M. Lifshitz, Oscillatory approach to a singular point in relativistic cosmology, *Adv. Phys.*, 19, 525, 1970
 - [12] C.W. Misner, Mixmaster Universe, *Phys. Rev. Lett.*, 22, 1071, 1969
 - [13] C.W. Misner, Quantum Cosmology, *Physical Review* 186, 5.1 319, 1969
 - [14] M.P. Ryan, Qualitative cosmology : Diagramatic solutions for Bianchi Type IX Universes with expansion, Rotation and Shear. I the symmetric case, *Ann. Phys.*, 65, 506, 1971
 - [15] R.T. Jantzen, Spatially homogeneous dynamics : A unified picture, version reformartée et corrigée de *Proc. Int. Sch. Phys. E. Fermi, Course LXXXVI (1982)*, on "Gamov Cosmology", R. Ruffini, F. Melchiorri Eds., North Holland, Amsterdam, 61-147, 1987 [*arXiv : gr - qc/0102035*]
 - [16] V.R. Gavrilov, V.D. Ivashchuk and V.N. Melnikov, Integrable pseudo-Euclidean Toda-like systems in multidimensional cosmology with multicomponent perfect fluid, *J. Math. Phys.*, 36, 5829, 1995
 - [17] A.A. Kirillov and V.N. Melnikov, Dynamics of inhomogeneities of the metric in the vicinity of a singularity in multidimensional cosmology, *Phys. Rev. D*, 52, 723, 1995
 - [18] A. E. Pavlov, The Mixmaster Cosmological Model as a pseudo-Euclidean generalized Toda chain, *Regul. & Chaotic Dyn.*, 1 (1), 111-119, 1996
 - [19] M. Szydlowski and M. Biesiada, Kovalewski exponents and integrability properties in class A homogeneous cosmological models, *J. non lin. math. phys.*, 9, L1, 2002
 - [20] R. Conte, The Painlevé approach to non linear ODE, p.77-180, in *The Painlevé property : One century later*, CRM series in Mathematical Physics, Editeur R. Conte, Springer Verlag, 1999
 - [21] H. Yoshida, Necessary condition for the existence of algebraic first integrals I : Kovalevski exponents, *Celestial mechanics*, 31, 363, 1983
 - [22] H. Yoshida, Necessary condition for the existence of algebraic first integrals II : Conditions for algebraic , *Celestial mechanics*, 31, 381, 1983
 - [23] Y. Elskens and M. Henneaux, Chaos in Kaluza-Klein models, *Class. Quantum Grav.*, 4, L161, 1987